



IN THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

1. (Currently Amended): A cosmetic composition comprising, in a physiologically acceptable medium, at least two dyes, including at least one photochromic dye and at least one goniochromatic coloring agent, wherein the photochromic dye has a difference in hue  $\Delta E$  between its excited state and its unexcited state at least equal to 5 and is at least one naphthopyran derivative compound of formula (Ia) or (IIa) below:



(Ia)



(IIa)

in which:

- R<sub>1</sub> represents:
  - (i) a hydrogen atom;
  - (ii) a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing from 1 to 30 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P, and/or optionally halogenated or perhalogenated;
  - (iii) a hydrocarbon-based ring formed with one of the bonds “f” or “gh” and the radical R<sub>7</sub>; or

- (iv) a group selected from the group consisting of  $-\text{COOR}_4$ ,  $-\text{C}(\text{O})\text{NR}_2\text{R}_3$ ,  $-\text{NR}_2\text{R}_3$ ,  $-\text{OR}_4$  and  $-\text{SR}_4$ , in which:


-  $\text{R}_2$  and  $\text{R}_3$  either represent, independently of each other, a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 1 to 20 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P,

or, taken together with the nitrogen atom to which they are attached, form a saturated or unsaturated hydrocarbon-based heterocycle containing 3 to 10 carbon atoms and optionally 1 to 5 other hetero atoms selected from the group consisting of N, O, S, Si and P, the ring optionally being substituted with at least one linear, branched or cyclic, saturated or unsaturated hydrocarbon-based radical containing 1 to 20 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P;


-  $\text{R}_4$  represents a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 1 to 20 carbon atoms, which is optionally halogenated or perhalogenated and/or optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P;

-  $\text{R}_5$  and  $\text{R}_6$  represent, independently of each other, a group selected from the group consisting of:

- (i) the saturated cyclic aminoaryl groups of formula (IIA) or (IIB):




(IIA)



(IIB)

in which the ring comprising N and X is a saturated ring containing in total 3 to 30 atoms including the nitrogen, the remainder being carbon atoms and/or hetero atoms selected from the group consisting of O, S, Si and P and/or groups selected from the group consisting of -NH and -NR with R representing a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based radical containing 1 to 20 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P;


- (ii) the indolinoaryl groups of formula (III):



(III)


in which  $R_{10}$  and  $R_{11}$  represent, independently of each other, a group selected from the group consisting of (i) linear, branched or cyclic, saturated or unsaturated hydrocarbon-based groups containing 1 to 30 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P, and/or optionally halogenated or perhalogenated; (ii) halogen atoms; (iii) -CN (nitrile), -COOH (carboxylate) or -NO<sub>2</sub> (nitro) groups; (iv) a hydrogen atom; (v) a group selected from the group consisting of -C(O)NR<sub>2</sub>R<sub>3</sub>, -NR<sub>2</sub>R<sub>3</sub>, -OR<sub>4</sub> and -SR<sub>4</sub> with R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> having the meanings given above; (vi) the radicals  $R_{10}$  and  $R_{11}$  together possibly forming a saturated or unsaturated hydrocarbon-based ring containing in total 5 to 8 atoms (including the atoms of the indoline ring), the atoms being selected from the group consisting of C, O, S and/or NR with R representing H or a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based radical containing 1 to 20 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P,

- (iii) the groups of formula (IV):




in which m and p are, independently of each other, integers ranging from 2 to 5;


- (iv) the unsaturated cyclic aminoaryl groups of formula (VA), (VB) or (VC):



(VA)



(VB)



(VC)

in which R<sub>8</sub> and R<sub>9</sub> represent, independently of each other, a group selected from the group consisting of (i) linear, branched or cyclic, saturated or unsaturated hydrocarbon-based groups containing 1 to 30 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P, and/or optionally halogenated or perhalogenated; (ii) halogen atoms; (iii) -CN (nitrile), -COOH (carboxylate) or -NO<sub>2</sub> (nitro) groups; (iv) a hydrogen atom; (v) a group selected from the group consisting of -C(O)NR<sub>2</sub>R<sub>3</sub>, -NR<sub>2</sub>R<sub>3</sub>, -OR<sub>4</sub> and -SR<sub>4</sub> with R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub> having the meanings given above;

- (v) a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 1 to 30 carbon atoms, optionally comprising 1 to 5 hetero atoms selected

from the group consisting of N, O, S, Si and P; and especially a group selected from the group consisting of  $-C_6H_4-CONR_2R_3$ ,  $-C_6H_4-NR_2R_3$  and  $-C_6H_4-OR_4$  with  $R_2$ ,  $R_3$  and  $R_4$  having the meanings given above;

- $R_7$  represents a group selected from the group consisting of:
  - (i) linear, branched or cyclic, saturated or unsaturated hydrocarbon-based groups containing 1 to 30 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P, and/or optionally halogenated or perhalogenated;
  - (ii) halogen atoms;
  - (iii)  $-CN$  (nitrile),  $-COOH$  (carboxylate),  $-NO_2$  (nitro),  $-N=N-$  (azo),  $=NH$  (imino) or  $-CONH_2$  (amide) groups;
  - (iv) a hydrogen atom;
  - (v) a group selected from the group consisting of  $-C(O)NR_2R_3$ ,  $-NR_2R_3$ ,  $-OR_4$  and  $-SR_4$  with  $R_2$ ,  $R_3$  and  $R_4$  having the meanings given above;
  - (vi) the radical  $R_7$  also possibly forming, with one of the bonds “i”, “j”, “k” or “g,h” taken with the radical  $R_1$ , or “f” taken with the radical  $R_1$ , a saturated hydrocarbon-based ring containing in total 3 to 8 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P;

- $R'_1$  represents a group selected from the group consisting of:
  - (i) a hydrogen atom;
  - (ii) a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 1 to 30 carbon atoms, optionally comprising 1 to 5 hetero atoms selected

from the group consisting of N, O, S, Si and P, and/or optionally halogenated or perhalogenated;

- (iii) a group selected from the group consisting of  $-C(O)NR_2R_3$ ,  $-NR_2R_3$ ,  $-OR_4$  and  $-SR_4$ , with  $R_2$ ,  $R_3$  and  $R_4$  having the meanings given above;

- $R'_2$  represents a group selected from the group consisting of:

- (i) linear, branched or cyclic, saturated or unsaturated hydrocarbon-based groups containing 1 to 30 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P, and/or optionally halogenated or perhalogenated;

- (ii) halogen atoms;

- (iii)  $-CN$  (nitrile),  $-COOH$  (carboxylate),  $-NO_2$  (nitro),  $-N=N-$  (azo),  $=NH$  (imino) or  $-CONH_2$  (amide) groups;

- (iv) a hydrogen atom;

- (v) a group selected from the group consisting of  $-C(O)NR_2R_3$ ,  $-NR_2R_3$ ,  $-OR_4$  and  $-SR_4$ , with  $R_2$ ,  $R_3$  and  $R_4$  having the meanings given above,

wherein the naphthopyran derivative is present in the composition in a dissolved form.

2. (Original): The composition according to claim 1, wherein the photochromic dye is present in an amount that is effective to give the composition a dynamic effect in terms of color and radiance.

3-4. (Canceled).

5. (Original): The composition according to claim 1, wherein  $R_1$  represents a hydrogen atom; a hydrocarbon-based ring with one of the bonds "f" or "gh" and the radical

R<sub>7</sub>; or a group selected from the group consisting of -COOR<sub>4</sub>, -NR<sub>2</sub>R<sub>3</sub>, -OR<sub>4</sub> and -SR<sub>4</sub>, in which:

- R<sub>2</sub> and R<sub>3</sub> either may represent, independently of each other, a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 1 to 20, carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P,

or, taken together with the nitrogen atom to which they are attached, may form a saturated or unsaturated hydrocarbon-based heterocycle containing 3 to 10 carbon atoms and optionally 1 to 5 other hetero atoms selected from the group consisting of N, O, S, Si and P, the ring optionally being substituted with at least one linear, branched or cyclic, saturated or unsaturated hydrocarbon-based radical containing 1 to 20 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P;


- R<sub>4</sub> may represent a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 1 to 20 carbon atoms, optionally halogenated or perhalogenated and/or optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P;

and/or


- R<sub>5</sub> and R<sub>6</sub> may represent, independently of each other, a group selected from the group consisting of:

- the saturated cyclic aminoaryl groups of formula (IIA) or (IIB):





(IIA)



(IIB)

in which the ring comprising N and X is a saturated ring which contains in total 3 to 30 including nitrogen, the rest being carbon atoms and/or hetero atoms selected from the group consisting of O, S, Si and P and/or groups selected from the group consisting of -NH and -NR with R representing a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based radical containing 1 to 20 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P;


- a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 1 to 30 carbon atoms, optionally comprising 1 to 5 hetero atoms selected from the group consisting of N, O, S, Si and P; and especially a group selected from the group consisting of  $-C_6H_4-CONR_2R_3$ ,  $-C_6H_4-NR_2R_3$  and  $-C_6H_4-OR_4$  with  $R_2$ ,  $R_3$  and  $R_4$  having the meanings given above.

6. (Canceled).

7. (Withdrawn): The composition according to claim 1, wherein the photochromic dye corresponds to formula (II) for which:

-  $R'_1$  represents hydrogen or a group  $-COOR$  with R being a saturated hydrocarbon-based radical containing 1 to 12 carbon atoms;  
and/or

-  $R_5$  and  $R_6$  represent, independently of each other, either (i) a group of formula (IIA):



in which the ring comprising N and X is a saturated ring containing in total 4 to 7 atoms, including nitrogen,

or (ii) a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 5 to 14 carbon atoms, optionally comprising 1 or 2 hetero atoms selected from the group consisting of N, O and S;

and/or

-  $R'_2$  represents hydrogen or a group  $-NR'R''$ , with  $R'$  and  $R''$ , which may be identical or different, representing a linear or branched, saturated hydrocarbon-based group containing 1 to 12 carbon atoms.

8. (Withdrawn): The composition according to claim 1, wherein the photochromic dye is selected from the group consisting of :

- 3,3-di(4-methoxyphenyl)-6-morpholino-3H-naphtho[2,1-b]pyran

- 3-phenyl-3-(4-morpholinophenyl)-6-morpholino-3H-naphtho[2,1-b]pyran
- 3-phenyl-3-(4-piperidinophenyl)-6-morpholino-3H-naphtho[2,1-b]pyran
- 3-phenyl-3-(4-piperidinophenyl)-6-carboxymethyl-9-N-dimethyl-3H-naphtho[2,1-b]pyran
- 2-phenyl-2-(4-piperidinophenyl)-5-carboxymethyl-9-N-dimethyl-2H-naphtho[1,2-b]pyran, and mixtures thereof.

9. (Withdrawn): The composition according to claim 1, comprising from 0.001% to 20% by weight of the photochromic dye, relative to the total weight of the cosmetic composition.

10. (Original): The composition according to claim 1, wherein the photochromic dye comprises at least two different naphthopyran derivatives.

11. (Canceled).

12. (Previously Presented): The composition according to claim 1, wherein the second dye is selected from the group consisting of the group consisting of goniochromatic coloring agents, water-soluble or liposoluble monochromatic dyes, pigments, reflective particles and nacres, and mixtures thereof.

13. (Canceled):

14. (Previously Presented): The composition according to claim 13, comprising from 0.1% to 60% of the goniochromatic coloring agent, relative to the total weight of the composition.

15. (Original): The composition according to claim 13, wherein the goniochromatic coloring agent is selected from the group consisting of multilayer interference structures and liquid-crystal coloring agents.

16. (Original): The composition according to claim 15, wherein the goniochromatic coloring agent of multilayer interference structure comprises at least two layers, each layer, optionally independently of the other layer(s), being made from at least one material selected from the group consisting of the group consisting of the following materials:  $\text{MgF}_2$ ,  $\text{CeF}_3$ ,  $\text{ZnS}$ ,  $\text{ZnSe}$ ,  $\text{Si}$ ,  $\text{SiO}_2$ ,  $\text{Ge}$ ,  $\text{Te}$ ,  $\text{Fe}_2\text{O}_3$ ,  $\text{Pt}$ ,  $\text{Va}$ ,  $\text{Al}_2\text{O}_3$ ,  $\text{MgO}$ ,  $\text{Y}_2\text{O}_3$ ,  $\text{S}_2\text{O}_3$ ,  $\text{SiO}$ ,  $\text{HfO}_2$ ,  $\text{ZrO}_2$ ,  $\text{CeO}_2$ ,  $\text{Nb}_2\text{O}_5$ ,  $\text{Ta}_2\text{O}_5$ ,  $\text{TiO}_2$ ,  $\text{Ag}$ ,  $\text{Al}$ ,  $\text{Au}$ ,  $\text{Cu}$ ,  $\text{Rb}$ ,  $\text{Ti}$ ,  $\text{Ta}$ ,  $\text{W}$ ,  $\text{Zn}$ ,  $\text{MoS}_2$ , cryolite, alloys and polymers, and combinations thereof.

17. (Withdrawn): The composition according to claim 15, comprising a liquid-crystal coloring agent obtained by polymerization of a monomer mixture comprising:

- a) at least a first monomer A of formula (I)  $\text{Y1-A1-M1-A2-Y2}$

in which

- i)  $\text{Y1}$  and  $\text{Y2}$ , which may be identical or different, represent an acrylate or methacrylate group;

- ii)  $\text{A1}$  and  $\text{A2}$ , which may be identical or different, represent a group of formula  $\text{-C}_n\text{H}_{2n}\text{-}$ , in which  $n$  is an integer ranging from 1 to 20;

- iii)  $\text{M1}$  denotes a group of general formula (I')  $\text{-R}_1\text{-X}_1\text{-R}_2\text{-X}_2\text{-R}_3\text{-X}_3\text{-R}_4\text{-}$ , in which  $\text{R}_1$  and  $\text{R}_4$  denote  $\text{-O-}$ , and  $\text{R}_2$  and  $\text{R}_3$  denote  $\text{-COO-}$ ,

and  $\text{X}_1$ ,  $\text{X}_2$  and  $\text{X}_3$  are a 1,4-phenylene group, the carbonyl group  $\text{-CO-}$  of  $\text{R}_2$  and of  $\text{R}_3$ , respectively, being linked to the group  $\text{X}_1$  or  $\text{X}_3$ , respectively,

and

- b) at least a second monomer B, which is chiral, of formula (II) V1-W1-Z-W2-V2,

in which

- i) V1 denotes an acrylate or methacrylate group, and V2 denotes a C<sub>1</sub>-C<sub>20</sub> alkyl,


C<sub>1</sub>-C<sub>20</sub> alkoxy, (C<sub>1</sub>-C<sub>20</sub>)alkoxycarbonyl or -OH group,;

- ii) W1 represents a divalent group of formula -X'1-CO-O-,

W2 represents a divalent group of formula -O-CO-X'1-,

in which X'1 denotes a 1,4-phenylene group,

and Z denotes a chiral group containing two bonds, derived from the dianhydrohexite group, in particular a divalent radical of formula:



18. (Withdrawn): Composition according to claim 17, wherein the liquid-crystal coloring agent is obtained from a monomer mixture comprising from 70% to 99% by weight of monomer A and from 1% to 30% by weight of monomer B, relative to the total weight of monomer A and of monomer B.

19. (Withdrawn): The composition according to claim 15, comprising a liquid-crystal coloring agent present in a content ranging from 0.01% to 99% by weight, relative to the total weight of the composition.

20. (Original): The composition according to claim 1, comprising, as a second dye, from 0.001% to 15% by weight of at least one liposoluble or water-soluble dye relative to the total weight of the composition.

21. (Original): The composition according to claim 1, comprising, as a second dye, from 0.01% to 25% by weight of at least one pigment relative to the total weight of the composition.

22. (Original): The composition according to claim 1, comprising, as a second dye, from 0.01% to 20% by weight of naces relative to the total weight of the composition.

23. (Original): The composition according to claim 1, comprising, as a second dye, from 0.1% to 20% of reflective particles relative to the total weight of the composition.

24. (Original): The composition according to claim 1, further comprising at least one oily phase.

25. (Original): The composition according to claim 24, wherein the oily phase comprises one or more polar or apolar, volatile or non-volatile oils.

26. (Original): The composition according to claim 24, wherein the oily phase comprises from 5% to 100% by weight, of polar oils relative to the total weight of the oily phase.

27. (Original): The composition according to claim 24, wherein the oily phase comprises from 5% to 100% by weight, of apolar oils relative to the total weight of the oily phase.

28. (Original): The composition according to claims 24, wherein the oils may be selected from the group consisting of volatile or non-volatile oils of animal, plant, mineral or synthetic origin, and mixtures thereof.

29. (Previously Presented): The composition according to claim 28, wherein the oils are selected from the group consisting of animal or plant oils, synthetic esters and ethers,

fatty acids, fatty alcohols, linear or branched hydrocarbons of mineral or synthetic origin, and glycerides, and mixtures thereof.

30. (Original): The composition according to claims 24, wherein the oily phase is such that the photochromic dye(s) is(are) soluble or dispersible therein.

31. (Original): The composition according to claims 24, wherein comprising from 1% to 99% by weight, of oily phase relative to the total weight of the composition.

32. (Original): The composition according to claim 1, comprising from 0.1% to 50% by weight, of a fatty substance other than an oil, relative to the total weight of the composition.

33. (Original): The composition according to claim 1, wherein said composition is anhydrous.

34. (Original): The composition according to claim 1, further comprising at least one aqueous phase.

35. (Original): The composition according to claim 34, wherein the aqueous phase comprises from 0.1% to 14% by weight of a C<sub>2</sub>-C<sub>6</sub> monoalcohol, relative to the total weight of the aqueous phase.

36. (Original): The composition according to claim 1, further comprising at least one surfactant.

37. (Original): The composition according to claim 36, wherein said surfactant is present in an amount from 0.01% to 30% by weight relative to the total weight of the composition.

38. (Original): The composition according to claim 1, further comprising at least one thickener.

39. (Original): The composition according to claim 38, wherein said thickener is present in an amount from 0.01% to 6% by weight relative to the total weight of the composition.

40. (Original): The composition according to claim 1, further comprising at least one film-forming polymer.

41. (Original): The composition according to claim 40, wherein said film-forming polymer is present in an amount from 0.01% to 4% by weight relative to the total weight of the composition.

42. (Original): The composition according to claim 1, further comprising at least one filler.

43. (Previously Presented): The composition according to claim 42, wherein said filler is present in a proportion of from 0.01% to 60% by weight relative to the total weight of the composition.


44. (Withdrawn): A process for making up the skin, the lips and/or the integuments, comprising applying the composition according to claim 1 to the skin, the lips and/or the integuments.

45. (Withdrawn): The process of Claim 44, comprising applying said composition to the skin.


46. (Withdrawn): The process according to claim 45, wherein the photochromic dye is a naphthopyran.



47. (Previously Presented): A cosmetic composition comprising, in a physiologically acceptable medium, at least two dyes, including at least one photochromic dye, wherein the photochromic dye has a difference in hue  $\Delta E$  between its excited state and its unexcited state at least equal to 5 and is at least one naphthopyran derivative compound of formula (Ia):




(Ia)



(IIa)


for which:

-  $R_1$  represents hydrogen; or a group  $-COOR$  with  $R$  being a saturated hydrocarbon-based radical containing 1 to 12 carbon atoms,; or a group




and/or

-  $R_5$  and  $R_6$  represent, independently of each other, either (i) a group of formula (IIA):



in which the ring comprising N and X is a saturated ring containing in total 4 to 7 atoms, including nitrogen, and especially 3 to 5 carbon atoms and 0 or 1 oxygen atom; and in particular a group of formula:



or (ii) a linear, branched or cyclic, saturated or unsaturated hydrocarbon-based group containing 5 to 14 carbon atoms, optionally comprising 1 or 2 hetero atoms selected from the group consisting of N, O or S;

and/or

$R_7$  represents a hydrogen atom or a group  $-NR_2R_3$ , with  $R_2$  and  $R_3$  representing, independently of each other, a linear or branched, saturated hydrocarbon-based group containing 1 to 12 carbon atoms.